



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Summing the series in parentheses we obtain

$$S = a \left\{ kn^2 - 2nk(k-1) + \frac{4k^3 - 6k^2 + 2k}{3} \right\} + b \left\{ kn^2 - 2nk^2 + \frac{4k^3 - k}{3} \right\},$$

which may be written as

$$(a+b)kn^2 - 2nk(ak+bk-a) + \left\{ \frac{4}{3}k^3(a+b) - 2ak^2 + \frac{k}{3}(2a-b) \right\}.$$

Also solved by H. C. FEEMSTER, HORACE OLSON and the PROPOSER.

**441. Proposed by W. D. CAIRNS, Oberlin College.**

Prove that the equation  $(e-1)x = e^x - 1$  has two and only two real roots.

### I. SOLUTION BY H. S. UHLER, Yale University.

Let  $y = e^x - 1 - (e-1)x$  and observe that  $y = 0$  for  $x = 0$  and  $x = 1$ . It remains to show that there can be no more real roots.

$$\frac{dy}{dx} = e^x - e + 1, \quad (1) \quad \frac{d^2y}{dx^2} = e^x. \quad (2)$$

Equation (1) shows that the slope of the tangent is positive for all values of  $x$  greater than  $\log_e(e-1)$  and negative for all values of  $x$  less than this value ( $x_0 = 0.541325$ ). Since  $e^x$  is essentially positive, equation (2) indicates formally that the single stationary point, indicated by equation (1), corresponds to a minimum value of  $y$ . The coördinates of the minimum are  $x_0 = \log_e(e-1)$  and  $y_0 = (e-1)(1-x_0) - 1 = -0.211867$ . It is clear, therefore, from the properties of the graph that the curve cannot cut the axis of  $x$  in more than two points and hence the given equation has two and only two real roots.

### II. SOLUTION BY GRACE M. BAREIS, Ohio State University.

Writing  $e$  in series form in each member and transposing, the equation becomes

$$x \left[ \frac{x-1}{2} + \frac{x^2-1}{3} + \frac{x^3-1}{4} + \dots \right] = 0, \text{ or } x(x-1) \left[ \frac{1}{2} + \frac{x+1}{3} + \frac{x^2+x+1}{4} + \dots \right] = 0.$$

Hence,  $x = 0$ ,  $x - 1 = 0$ , or  $\frac{1}{2} + \frac{x+1}{3} + \frac{x^2+x+1}{4} + \dots = 0$ . Since each term of the left member of the last equation is positive, this equation can have no real positive root.

To prove that the original equation can have no real negative root, put it in the form

$$x = \frac{e^x - 1}{e - 1},$$

or  $x = e^{x-1} + e^{x-2} + e^{x-3} + \dots$ , a convergent series. Any real negative number makes the left member negative but the right member positive, and hence there are no real negative roots. Hence, 0 and 1 are the only real roots.

Solutions were also received from W. L. AGARD, F. L. GRIFFIN, H. C. FEEMSTER, WALTER C. EELLS, HORACE OLSON, C. E. HORNE, IRBY C. NICHOLS, and FRANK IRWIN.

### GEOMETRY.

**467. Proposed by E. T. BELL, Seattle, Washington.**

It is well-known that if  $i, j, k, l$  are concyclic points,  $W_i$  the Wallace line (frequently, and erroneously, called the Simson line), of  $i$  with respect to the triangle  $jkl$ , then  $W_i, W_j, W_k, W_l$  are concurrent, say in the point  $\{i, j, k, l\}$ . If 1, 2, 3,  $\dots$  denote concyclic points, prove that:

(i)  $\{1, 2, 3, 4\}$ ,  $\{1, 2, 3, 5\}$ ,  $\{1, 2, 4, 5\}$ ,  $\{1, 3, 4, 5\}$ ,  $\{2, 3, 4, 5\}$  are concyclic; say on the circle [1, 2, 3, 4, 5];

(ii) Starting with 1, 2, 3, 4, 5, 6, omitting each point in turn, by (i), six circles, are found; these are concurrent, say in the point  $\{1, 2, 3, 4, 5, 6\}$ ;

(iii) Starting with 1, 2, 3, 4, 5, 6, 7, seven points of the kind in (ii) are found; these lie on a circle.

(iv) Continuing thus indefinitely, there is, in each case, finally a unique point or circle according as the number of initial points is even or odd. Also, at any stage, the point of concurrence on the circle bears a simple relation to the initial points: what is it?

## I. SOLUTION BY J. W. CLAWSON, Collegeville, Pa.

If  $H_i$  is the orthocenter of triangle  $jkl$ ,  $O$  the center of the circle, radius  $R$ , on which the points  $i, j, k, l$  lie,—

$iH_j$  = twice distance of  $O$  from  $kl = jH_i$ . Also  $iH_j$  is parallel to  $jH_i$ ; hence  $H_iH_j$  is equal to, and parallel to,  $ij$ . It follows that  $H_iH_jH_kH_l$  is inversely congruent with  $ijkl$ , the corresponding sides of the quadrilaterals being equal and parallel, but arranged in opposite orders.

Again,  $iH_i, jH_j, kH_k, lH_l$  have a common mid-point; and, since  $iH_i$  bisects  $W_i$  and so on, this common point is the point of concurrency of  $W_i, W_j, W_k, W_l$ . The point of concurrency  $\{i, j, k, l\}$  bisects the join of  $O$  to the center of the circle circumscribing  $H_iH_jH_kH_l$ .

(i) If  $H_{123}$  is the orthocenter of triangle 123,  $H_{123}, H_{234}, H_{341}, H_{412}$  are four vertices of a pentagon inversely congruent with 12345. Moreover the sides of this pentagon are parallel to the corresponding sides of the pentagon 12345. Call the center of the circumcircle of this pentagon,  $P_5$ . Similarly  $H_{123}, H_{235}, H_{351}, H_{512}$  and three other sets of orthocenters determine four other congruent pentagons with parallel sides. Each of the orthocenters occurs as a vertex of two of the pentagons. It follows that  $P_5, P_4, P_3, P_2, P_1$ , the centers of the circles circumscribing these five pentagons, form a pentagon which is congruent with each of the pentagons and hence inversely congruent with the original pentagon 12345. The points  $P_5, P_4, P_3, P_2, P_1$  are therefore concyclic. Call the center of their circle,  $O_6$ .

It follows at once that the points (1234), (2345), (3451), (4512), (5123) are concyclic on a circle of half the radius of the given circle, its center bisecting the join of  $O$  to  $O_6$ .

(ii) The circle, center  $O_6$ , passes through  $P_5$ , the center of a circle determined by the orthocenters of the triangles obtained by selecting triads of points from 1, 2, 3, 4. In the same way if we consider the five points 1, 2, 3, 4, 6 we shall obtain a circle, center  $O_5$ , which passes through  $P_5$  and through four new points. In this way we can get six hexagons, one of which has  $P_1, P_2, P_3, P_4, P_5$  for five of its vertices, each inversely congruent with the hexagon 123456 and five vertices of each hexagon (orthocenters of triangles obtained by selecting three points from the six) belonging also to a different hexagon of the system. It follows that  $O_6O_5O_4O_3O_2O_1$  forms a hexagon which is congruent with each of the hexagons and hence is inversely congruent with the original hexagon 123456. Hence  $O_6, O_5, O_4, O_3, O_2, O_1$  are concyclic; and the circles with  $O_6, O_5, O_4, O_3, O_2, O_1$  as centers and with  $R$  as radius concur at a point, say  $Q_7$ .

It follows at once that circles (1, 2, 3, 4, 5),  $\dots$  concur at a point halfway between  $Q_7$  and  $O$ .

(iii) This process can be continued indefinitely as stated in the problem.

## II. SOLUTION BY NORMAN ANNING, Chilliwick, B. C.

Letting  $e^{i\phi} = \cos \phi + i \sin \phi$ , then  $Re^{i\theta_j}$  ( $j = 1, 2, 3, 4 \dots$ ) are points on a circle of radius  $R$ , whose center is at the origin. The  $W$  line of 4 qua 123 passes through the foot of the perpendicular from 4 on line 12 and makes with this perpendicular an angle equal to that between 13 and 14. All points on  $W_4$  are given by

$$Re^{i\theta_4} + ke^{i\frac{\theta_1+\theta_2}{2}} + xe^{i\frac{\theta_1+\theta_2+\theta_3-\theta_4}{2}},$$

where  $x$  is a running coördinate on the line and  $k$ , the distance from 4 to 12, is to be determined. Projecting on the radius vector midway between 1 and 2,

$$k + R \cos \left( \theta_4 - \frac{\theta_1 + \theta_2}{2} \right) = R \cos \frac{\theta_1 - \theta_2}{2}, \quad k = 2R \sin \frac{\theta_4 - \theta_2}{2} \sin \frac{\theta_4 - \theta_1}{2}.$$

$$W_4 \text{ is } Re^{i\theta_4} + 2R \sin \frac{\theta_4 - \theta_2}{2} \sin \frac{\theta_4 - \theta_1}{2} e^{i\frac{\theta_1+\theta_2}{2}} + xe^{i\frac{\theta_1+\theta_2+\theta_3-\theta_4}{2}}.$$

$$W_3 \text{ is } Re^{i\theta_3} + 2R \sin \frac{\theta_3 - \theta_2}{2} \sin \frac{\theta_3 - \theta_1}{2} e^{i\frac{\theta_1+\theta_2}{2}} + ye^{i\frac{\theta_1+\theta_2-\theta_3+\theta_4}{2}}.$$

